

ON HYPERGEOMETRIC BERNOULLI NUMBERS AND POLYNOMIALS

SU HU AND MIN-SOO KIM

ABSTRACT. In this note, we shall provide several properties of hypergeometric Bernoulli numbers and polynomials, including sums of products identity, differential equations and recurrence formulas.

1. INTRODUCTION

For $N \in \mathbb{N}$, Howard [11, 12] defined the hypergeometric Bernoulli polynomials $B_{N,n}(x)$ by the generating function

$$(1.1) \quad \frac{t^N e^{xt}/N!}{e^t - T_{N-1}(t)} = \sum_{n=0}^{\infty} B_{N,n}(x) \frac{t^n}{n!},$$

where $T_{N-1}(t)$ is the Taylor polynomial of order $N-1$ for the exponential function. In particular, when $N=1$, we recover the classical Bernoulli polynomials, i.e. $B_{1,n}(x) = B_n(x)$ (see (1.2) below).

In this note, we shall give an expression for a sum of products of the hypergeometric Bernoulli polynomials (see Theorem 1.2 below), generalizing some classical results for Bernoulli polynomials. We also present a differential equation and a recurrence relation for these polynomials (see Theorems 1.5 and 1.8).

Recall that, the Bernoulli polynomials is defined by the generating function

$$(1.2) \quad \frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!},$$

while the Bernoulli numbers is defined by $B_n = B_n(0)$.

The Bernoulli numbers and polynomials have many applications and satisfy many interesting identities. The most remarkable one is Euler's sums of products identity

$$(1.3) \quad \sum_{i=0}^n \binom{n}{i} B_i B_{n-i} = -nB_{n-1} - (n-1)B_n \quad (n \geq 1).$$

This identity has been generalized by many authors from different directions (see [2, 4, 6, 14, 15, 20, 21]). In particular, Dilcher [6] provided explicit expressions for sums of products for arbitrarily many Bernoulli numbers and polynomials.

2000 *Mathematics Subject Classification.* Primary 11M35; Secondary 11B68.

Key words and phrases. Hypergeometric Bernoulli numbers and polynomials, Sums of products, Differential equations, Recurrence formulas, Appell polynomials.

Another approach to Bernoulli polynomials is to define them as an Appell sequence with zero mean:

$$(1.4) \quad B_0(x) = 1,$$

$$(1.5) \quad B'_n(x) = nB_{n-1}(x),$$

$$(1.6) \quad \int_0^1 B_n(x)dx = \begin{cases} 1 & n = 0 \\ 0 & n > 0. \end{cases}$$

Based on the properties of Appell polynomials, He and Ricci [10] proved that the Bernoulli polynomials $B_n(x)$ satisfy the following differential equations:

$$(1.7) \quad \frac{B_n}{n!}y^{(n)} + \frac{B_{n-1}}{(n-1)!}y^{(n-1)} + \cdots + \frac{B_2}{2!}y'' - \left(x - \frac{1}{2}\right)y' + ny = 0.$$

Howard [11, 12] gave a generalization of Bernoulli polynomials by considering the following generating function:

$$(1.8) \quad \frac{t^2 e^{xt}/2}{e^t - 1 - t} = \sum_{n=0}^{\infty} A_n(x) \frac{t^n}{n!}$$

and more generally, for all positive integer N

$$(1.9) \quad \frac{t^N e^{xt}/N!}{e^t - T_{N-1}(t)} = \sum_{n=0}^{\infty} B_{N,n}(x) \frac{t^n}{n!},$$

where $T_{N-1}(t)$ is the Taylor polynomial of order $N-1$ for the exponential function. For the cases $N=1$ and $N=2$, (1.9) reduces to (1.2) and (1.8), respectively. We see that the polynomials $B_{N,n}(x)$ have rational coefficients.

The polynomials $B_{N,n}(x)$ are named hypergeometric Bernoulli polynomials, while the numbers $B_{N,n} = B_{N,n}(0)$ are named hypergeometric Bernoulli numbers since the generating function $f(t) = \frac{e^t - T_{N-1}(t)}{t^N/N!}$ can be expressed as ${}_1F_1(1, N+1; t)$, where the confluent hypergeometric function ${}_1F_1(a, b; t)$ is defined by

$${}_1F_1(a, b; t) = \sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n} \frac{t^n}{n!}$$

and $(a)_n$ is the Pochhammer symbol

$$(a)_n = \begin{cases} a(a+1) \cdots (a+n-1) & (n \geq 1) \\ 1 & (n = 0) \end{cases}$$

(see [13, p. 2261]).

As their classical counterparts, the Bernoulli numbers and polynomials, the generalized Bernoulli numbers and polynomials, the hypergeometric Bernoulli numbers and polynomials also satisfy many interesting properties ([3, 7, 8, 16, 18, 19]).

Kamano [13] proved the following result for sums of products of hypergeometric Bernoulli numbers, which is a generalization of the works by Euler and Dilcher (see [6]).

Theorem 1.1 (Kamano [13, p. 2262, Main Theorem]). *Let N and r be positive integers. For any integer $n \geq r - 1$, we have*

$$\begin{aligned} \sum_{\substack{i_1, \dots, i_r \geq 0 \\ i_1 + \dots + i_r = n}} \frac{n!}{i_1! \dots i_r!} B_{N, i_1} \dots B_{N, i_r} \\ = \frac{1}{N^{r-1}} \sum_{i=0}^{r-1} A_r^{(N)}(i; 1 + N(r-1) - n) (-1)^i \binom{n}{i} i! B_{N, n-i}, \end{aligned}$$

where $A_r^{(N)}(i; s) \in \mathbb{Q}[s]$ ($0 \leq i \leq r-1$) are polynomials defined by the following recurrence relation:

$$(1.10) \quad \begin{aligned} A_1^{(N)}(0; s) &= 1 \\ A_r^{(N)}(i; s) &= \frac{s-1}{r-1} A_{r-1}^{(N)}(i; s-N) + A_{r-1}^{(N)}(i-1; s-N+1). \end{aligned}$$

Here $r \geq 2$ and $A_r^{(N)}(i; s)$ are defined to be zero for $i \leq -1$ and $i \geq r$.

As for Bernoulli polynomials, another approach to hypergeometric Bernoulli polynomials is to define them in terms of Appell sequence with zero mean (comparing with (1.4), (1.5) and (1.6) above):

$$(1.11) \quad B_{N,0}(x) = 1,$$

$$(1.12) \quad B'_{N,n}(x) = n B_{N,n-1}(x),$$

$$(1.13) \quad \int_0^1 (1-x)^{N-1} B_{N,n}(x) dx = \begin{cases} \frac{1}{N} & n = 0 \\ 0 & n > 0 \end{cases}$$

(see [7, p. 768]).

This paper contains further properties of the hypergeometric Bernoulli numbers and polynomials, including a generalization of Kamano's result.

Theorem 1.2. *Let N and r be positive integers and let $x = x_1 + \dots + x_r$. For any integer $n \geq r - 1$, we have*

$$\begin{aligned} \sum_{\substack{i_1, \dots, i_r \geq 0 \\ i_1 + \dots + i_r = n}} \frac{n!}{i_1! \dots i_r!} B_{N, i_1}(x_1) \dots B_{N, i_r}(x_r) \\ = \frac{1}{N^{r-1}} \sum_{i=0}^{r-1} A_r^{(N)}(i, x; 1 + N(r-1) - n) (-1)^i \binom{n}{i} i! B_{N, n-i}(x), \end{aligned}$$

where $A_r^{(N)}(i, x; s) \in \mathbb{Q}[x, s]$ ($0 \leq i \leq r-1$) are polynomials defined by the recurrence relation:

$$(1.14) \quad \begin{aligned} A_1^{(N)}(0, x; s) &= 1 \\ A_r^{(N)}(i, x; s) &= \frac{s-1}{r-1} A_{r-1}^{(N)}(i, x; s-N) - \frac{x-(r-1)}{r-1} A_{r-1}^{(N)}(i-1, x; s-N+1). \end{aligned}$$

Here $r \geq 2$ and $A_r^{(N)}(i, x; s)$ are defined to be zero for $i \leq -1$ and $i \geq r$.

Remark 1.3. Theorem 1.1 is the special case if $x = 0$. Nguyen and Cheong [19] also obtained a similar statement using a result by Nguyen [18].

Remark 1.4. Let $r = 2, 3$ in (1.14) to obtain

$$\begin{aligned}
 (1.15) \quad & A_2^{(N)}(0, x; 1 + N - n) = N - n, \\
 & A_2^{(N)}(1, x; 1 + N - n) = -(x - 1), \\
 & A_3^{(N)}(0, x; 1 + 2N - n) = \frac{1}{2}(2N - n)(N - n), \\
 & A_3^{(N)}(1, x; 1 + 2N - n) = -\frac{1}{2}(2N - n)(x - 1) - \frac{1}{2}(x - 2)(N - n + 1), \\
 & A_3^{(N)}(2, x; 1 + 2N - n) = \frac{1}{2}(x - 2)(x - 1),
 \end{aligned}$$

thus

$$\begin{aligned}
 (1.16) \quad & \sum_{\substack{i_1, i_2 \geq 0 \\ i_1 + i_2 = n}} \frac{n!}{i_1! i_2!} B_{N, i_1}(x_1) B_{N, i_2}(x_2) \\
 &= \frac{1}{N}(N - n) B_{N, n}(x) + \frac{n}{N}(x - 1) B_{N, n-1}(x) \\
 &\quad (\text{where } x = x_1 + x_2 \text{ and } n \geq 1),
 \end{aligned}$$

and

$$\begin{aligned}
 (1.17) \quad & \sum_{\substack{i_1, i_2, i_3 \geq 0 \\ i_1 + i_2 + i_3 = n}} \frac{n!}{i_1! i_2! i_3!} B_{N, i_1}(x_1) B_{N, i_2}(x_2) B_{N, i_3}(x_3) \\
 &= \frac{1}{2N^2} \left[(N - n)(2N - n) B_{N, n}(x) \right. \\
 &\quad + n((2N - n)(x - 1) + (x - 2)(N - n + 1)) B_{N, n-1}(x) \\
 &\quad \left. + n(n - 1)(x - 1)(x - 2) B_{N, n-2}(x) \right] \\
 &\quad (\text{where } x = x_1 + x_2 + x_3 \text{ and } n \geq 2).
 \end{aligned}$$

Nguyen and Cheong [19, Example 18] also obtained (1.16) and (1.17). If $N = 1$, $x = 0$ and $r = 2$, then (1.16) becomes (1.3).

For $N, r \in \mathbb{N}$, the higher order hypergeometric Bernoulli polynomials $B_{N, n}^{(r)}(x)$ are defined by the generating function

$$(1.18) \quad \left(\frac{t^N / N!}{e^t - T_{N-1}(t)} \right)^r e^{xt} = \sum_{n=0}^{\infty} B_{N, n}^{(r)}(x) \frac{t^n}{n!}.$$

The higher order hypergeometric Bernoulli numbers are defined by $B_{N, n}^{(r)} = B_{N, n}^{(r)}(0)$ (see [13, 19]). The hypergeometric Bernoulli case corresponds to the special value $r = 1$. In particular, $B_{N, n}^{(1)}(x) = B_{N, n}(x)$, the hypergeometric Bernoulli polynomials, and $B_{N, n}^{(1)}(0) = B_{N, n}$, the hypergeometric Bernoulli numbers.

If we put $N = 1$ in (1.18), then we have $B_{1,n}^{(r)}(x) = B_n^{(r)}(x)$ the classical higher order Bernoulli polynomials. The classical higher order Bernoulli numbers is defined by $B_n^{(r)} = B_n^{(r)}(0)$ (see [6, 14, 16, 21]).

Using results established in [10] for the classical Bernoulli polynomials a differential equation for the higher order Bernoulli polynomials is derived next. The proof is based on the properties of Appell polynomials [1, 22].

Theorem 1.5. *The higher order hypergeometric Bernoulli polynomials $B_{N,n}^{(r)}(x)$ satisfy the differential equation*

$$\frac{B_{N,n}}{n!}y^{(n)} + \frac{B_{N,n-1}}{(n-1)!}y^{(n-1)} + \cdots + \frac{B_{N,2}}{2!}y'' - \left(\frac{x}{rN} - \frac{1}{N(N+1)}\right)y' + \frac{n}{rN}y = 0.$$

Corollary 1.6 (Lu [16, Theorem 2.2]). *The classical higher order Bernoulli polynomials $B_n^{(r)}(x)$ satisfy the differential equation*

$$\frac{B_n(1)}{n!}y^{(n)} + \frac{B_{n-1}(1)}{(n-1)!}y^{(n-1)} + \cdots + \frac{B_2(1)}{2!}y'' - \left(\frac{x}{r} - \frac{1}{2}\right)y' + \frac{n}{r}y = 0.$$

Theorem 1.5 in the case $N = r = 1$ gives the next result.

Corollary 1.7 (He and Ricci [10, Theorem 2.3]). *The classical Bernoulli polynomials $B_n(x)$ satisfy the differential equation*

$$\frac{B_n}{n!}y^{(n)} + \frac{B_{n-1}}{(n-1)!}y^{(n-1)} + \cdots + \frac{B_2}{2!}y'' - \left(x - \frac{1}{2}\right)y' + ny = 0.$$

A linear recurrence for higher order Bernoulli polynomials generalizing results of Lu [16] appears as consequence of the proof.

Theorem 1.8. *For $n \in \mathbb{N}$, the higher order hypergeometric Bernoulli polynomials $B_{N,n}^{(r)}(x)$ satisfy the recurrence*

$$B_{N,n+1}^{(r)}(x) = \left(x - \frac{r}{N+1}\right) B_{N,n}^{(r)}(x) - rN \sum_{k=0}^{n-1} \binom{n}{k} \frac{B_{N,n-k+1}}{n-k+1} B_{N,k}^{(r)}(x).$$

The special case $N = 1$ gives the following statement.

Corollary 1.9 (Lu [16, Theorem 2.1]). *For $n \in \mathbb{N}$, the higher order Bernoulli polynomials $B_n^{(r)}(x)$ satisfy the recurrence*

$$B_{n+1}^{(r)}(x) = \left(x - \frac{1}{2}r\right) B_n^{(r)}(x) - r \sum_{k=0}^{n-1} \binom{n}{k} \frac{B_{n-k+1}(1)}{n-k+1} B_k^{(r)}(x).$$

Letting $N = r = 1$ and replace n by $n+1$ in Theorem 1.8, we obtain the next result.

Corollary 1.10 (He and Ricci [10, Theorem 2.2]). *For $n \in \mathbb{N}$, the Bernoulli polynomials $B_n(x)$ satisfy the recurrence*

$$B_n(x) = \left(x - \frac{1}{2}\right) B_{n-1}(x) - \frac{1}{n} \sum_{k=0}^{n-2} \binom{n}{k} B_{n-k} B_k(x).$$

Remark 1.11. The special case $r = 1$ gives some results presented in [17].

2. PROOF OF THEOREM 1.2

Introduce the notations

$$(2.1) \quad F_{r,N}(t, x) = \left(\frac{t^N/N!}{e^t - T_{N-1}(t)} \right)^r e^{xt},$$

$$(2.2) \quad F_{r,N}(t) = F_{r,N}(t, 0) = \left(\frac{t^N/N!}{e^t - T_{N-1}(t)} \right)^r$$

and

$$(2.3) \quad F_N(t) = F_{1,N}(t) = \frac{t^N/N!}{e^t - T_{N-1}(t)}.$$

The proof begins with an auxiliary result.

Lemma 2.1.

- (i) $\frac{d}{dt}F_N(t) = \frac{N}{t}F_N(t) - F_N(t) - \frac{N}{t}F_{2,N}(t).$
- (ii) $\frac{d}{dt}F_{r,N}(t, x) = \frac{rN}{t}F_{r,N}(t, x) + (x - r)F_{r,N}(t, x) - \frac{rN}{t}F_{r+1,N}(t, x).$

Proof. Differentiating (2.3) with respect to t , yields

$$\begin{aligned} \frac{d}{dt}F_N(t) &= \frac{Nt^{N-1}/N!(e^t - T_{N-1}(t)) - t^N/N!(e^t - T_{N-2}(t))}{(e^t - T_{N-1}(t))^2} \\ &= \frac{N}{t} \frac{t^N/N!}{(e^t - T_{N-1}(t))} - \frac{t^N/N!(e^t - T_{N-1}(t) + t^{N-1}/(N-1)!)}{(e^t - T_{N-1}(t))^2} \\ &= \frac{N}{t}F_N(t) - F_N(t) - \frac{N}{t}F_{2,N}(t), \end{aligned}$$

giving Part (i). Part (ii) follows from Part (i). $\square$

The notations introduced above states that

$$(2.4) \quad F_{r,N}(t, x) = \sum_{n=0}^{\infty} B_{N,n}^{(r)}(x) \frac{t^n}{n!}.$$

Comparing coefficients of equal powers in Lemma 2.1(ii) gives the recurrence stated next.

Lemma 2.2. *Let $n \in \mathbb{N}$. Then*

$$B_{N,n}^{(r+1)}(x) = \frac{1}{N} \left(N - \frac{n}{r} \right) B_{N,n}^{(r)}(x) + \frac{1}{N} \frac{n}{r} (x - r) B_{N,n-1}^{(r)}(x).$$

Let $\binom{n}{i_1, \dots, i_r} = \frac{n!}{i_1! \cdots i_r!}$ be the multinomial coefficient. The next result gives $B_{N,n}^{(r)}(x)$ in terms of $B_{N,i}(x)$.

Lemma 2.3. *The identity holds for $n, r, N \in \mathbb{N}$*

$$(2.5) \quad B_{N,n}^{(r)}(x) = \sum_{\substack{i_1 + \cdots + i_r = n \\ i_1, \dots, i_r \geq 0}} \binom{n}{i_1, \dots, i_r} B_{N,i_1}(x_1) \cdots B_{N,i_r}(x_r)$$

with $x = x_1 + \cdots + x_r$.

Proof of the Theorem 1.2. The proof proceeds by induction.

The case $r = 1$ is clear. Lemma 2.2 gives

$$\begin{aligned}
B_{N,n}^{(r)}(x) &= \frac{1}{N} \left(N - \frac{n}{r-1} \right) B_{N,n}^{(r-1)}(x) + \frac{1}{N} \frac{n}{r-1} (x - (r-1)) B_{N,n-1}^{(r-1)}(x) \\
&= \frac{1}{N^{r-1}} \left(N - \frac{n}{r-1} \right) \sum_{i=0}^{r-2} A_{r-1}^{(N)}(i, x; 1 + N(r-2) - n) \\
&\quad \times (-1)^i \binom{n}{i} i! B_{N,n-i}(x) \\
&\quad + \frac{1}{N^{r-1}} \frac{n}{r-1} (x - (r-1)) \sum_{i=0}^{r-2} A_{r-1}^{(N)}(i, x; 1 + N(r-2) - (n-1)) \\
&\quad \times (-1)^i \binom{n-1}{i} i! B_{N,n-1-i}(x) \\
&= \frac{1}{N^{r-1}} \left[\left(N - \frac{n}{r-1} \right) \sum_{i=0}^{r-2} A_{r-1}^{(N)}(i, x; 1 + N(r-2) - n) \right. \\
&\quad \times (-1)^i \binom{n}{i} i! B_{N,n-i}(x) \\
&\quad + \frac{n}{r-1} (x - (r-1)) \sum_{i=1}^{r-1} A_{r-1}^{(N)}(i-1, x; 1 + N(r-2) - (n-1)) \\
&\quad \times (-1)^{i-1} \binom{n-1}{i-1} (i-1)! B_{N,n-i}(x) \left. \right] \\
&= \frac{1}{N^{r-1}} \sum_{i=0}^{r-1} \left[\left(N - \frac{n}{r-1} \right) A_{r-1}^{(N)}(i, x; 1 + N(r-2) - n) (-1)^i \binom{n}{i} i! \right. \\
&\quad + \frac{n}{r-1} (x - (r-1)) A_{r-1}^{(N)}(i-1, x; 1 + N(r-2) - (n-1)) \\
&\quad \times (-1)^{i-1} \binom{n-1}{i-1} (i-1)! \left. \right] B_{N,n-i}(x) \\
&= \frac{1}{N^{r-1}} \sum_{i=0}^{r-1} \left[\frac{N(r-1) - n}{r-1} A_{r-1}^{(N)}(i, x; 1 + N(r-2) - n) \right. \\
&\quad - \frac{1}{r-1} (x - (r-1)) A_{r-1}^{(N)}(i-1, x; 1 + N(r-2) - (n-1)) \left. \right] \\
&\quad \times (-1)^i \binom{n}{i} i! B_{N,n-i}(x) \\
&= \frac{1}{N^{r-1}} \sum_{i=0}^{r-1} A_r^{(N)}(i, x; 1 + N(r-1) - n) (-1)^i \binom{n}{i} i! B_{N,n-i}(x) \\
&\quad (\text{using } s = 1 + N(r-1) - n \text{ in (1.14)}).
\end{aligned}$$

Lemma 2.3 gives the result. $\square$

3. DIFFERENTIAL EQUATIONS

3.1. Appell polynomials. The proof for the differential equation satisfied by the higher order hypergeometric Bernoulli polynomials requires some basic facts on Appell polynomials. These also appear in [10].

The Appell polynomials [1] are defined by the generating function:

$$(3.1) \quad A(t)e^{xt} = \sum_{n=0}^{\infty} \frac{R_n(x)}{n!} t^n,$$

where

$$(3.2) \quad A(t) = \sum_{n=0}^{\infty} \frac{R_n}{n!} t^n, \quad A(0) \neq 0$$

is an analytic function at $t = 0$, and $R_n = R_n(0)$.

A polynomial $p_n(x)$ ($n \in \mathbb{N}, x \in \mathbb{C}$) is said to be a quasi-monomial [16] if there are two operators, $\hat{M}, \hat{P}$, called the multiplicative and derivative operators, such that

$$(3.3) \quad \hat{M}(p_n(x)) = p_{n+1}(x)$$

and

$$(3.4) \quad \hat{P}(p_n(x)) = np_{n-1}(x),$$

where it is assumed (as usual) that

$$p_0(x) = 1 \quad \text{and} \quad p_{-1}(x) = 0.$$

The operators $\hat{M}$ and $\hat{P}$ must satisfy the commutation relation

$$(3.5) \quad [\hat{P}, \hat{M}] = \hat{P}\hat{M} - \hat{M}\hat{P} = \hat{I},$$

where $\hat{I}$ denotes the identity operator.

He and Ricci [10] showed that the multiplicative and derivative operators of $R_n(x)$ are

$$(3.6) \quad \hat{M} = (x + \alpha_0) + \sum_{k=0}^{n-1} \frac{\alpha_{n-k}}{(n-k)!} D_x^{n-k}$$

and

$$(3.7) \quad \hat{P} = D_x,$$

where $D_x = \frac{\partial}{\partial x}$ and the coefficients α_n are defined by

$$(3.8) \quad \frac{A'(t)}{A(t)} = \sum_{n=0}^{\infty} \alpha_n \frac{t^n}{n!}.$$

3.2. Proofs of Theorems 1.5 and 1.8. The proofs begin with an auxiliary result.

Lemma 3.1.

$$\frac{d^p}{dx^p} B_{N,n}^{(r)}(x) = \frac{n!}{(n-p)!} B_{N,n-p}^{(r)}(x), \quad n \geq p.$$

Proof. Start with

$$(3.9) \quad \left(\frac{t^N/N!}{e^t - T_{N-1}(t)} \right)^r \frac{d^p}{dx^p} e^{xt} = \left(\frac{t^N/N!}{e^t - T_{N-1}(t)} \right)^r (e^{xt} t^p) \\ = \left(\left(\frac{t^N/N!}{e^t - T_{N-1}(t)} \right)^r e^{xt} \right) t^p.$$

Substituting (1.18) into the right hand side of (3.9), we have

$$(3.10) \quad \left(\frac{t^N/N!}{e^t - T_{N-1}(t)} \right)^r \frac{d^p}{dx^p} e^{xt} = \left(\sum_{n=0}^{\infty} B_{N,n}^{(r)}(x) \frac{t^n}{n!} \right) t^p \\ = \sum_{n=0}^{\infty} B_{N,n}^{(r)}(x) \frac{t^{n+p}}{n!} \\ = \sum_{n=p}^{\infty} B_{N,n-p}^{(r)}(x) \frac{t^n}{(n-p)!} \\ = \sum_{n=p}^{\infty} \frac{B_{N,n-p}^{(r)}(x)}{(n-p)!} t^n.$$

Differentiating both sides of (1.18) with respect to x and using (3.10) gives

$$(3.11) \quad \sum_{n=p}^{\infty} \frac{B_{N,n-p}^{(r)}(x)}{(n-p)!} t^n = \sum_{n=0}^{\infty} \frac{d^p}{dx^p} \frac{B_{N,n}^{(r)}(x)}{n!} t^n.$$

Comparing the coefficients of both sides of the above equality, produces the result. $\square$

Proof of the Theorem 1.8. From (1.18), (3.1) and (3.2), we know that the hypergeometric Bernoulli polynomials $B_{N,n}^{(r)}(x)$ are Appell polynomials with

$$A(t) = \left(\frac{t^N/N!}{e^t - T_{N-1}(t)} \right)^r.$$

Using Lemma 2.1(ii) with $x = 0$ gives

$$\begin{aligned}
 \frac{A'(t)}{A(t)} &= \frac{rN}{t} - r - \frac{rN}{t} \left(\frac{t^N/N!}{e^t - T_{N-1}(t)} \right) \\
 &= \frac{rN}{t} \left(1 - \frac{t}{N} - \sum_{n=0}^{\infty} B_{N,n} \frac{t^N}{n!} \right) \\
 (3.12) \quad &= \frac{rN}{t} \left(-\frac{1}{N(N+1)}t - \sum_{n=2}^{\infty} B_{N,n} \frac{t^N}{n!} \right) \\
 &= r \left(-\frac{1}{N+1} - N \sum_{n=1}^{\infty} \frac{B_{N,n+1}}{n+1} \frac{t^N}{n!} \right).
 \end{aligned}$$

Here we use $B_{N,0} = 1$ and $B_{N,1} = -1/(N+1)$. Then (3.6)–(3.8) give the multiplicative and derivative operators of the hypergeometric Bernoulli polynomials:

$$(3.13) \quad \hat{M} = \left(x - \frac{r}{N+1} \right) - rN \sum_{k=0}^{n-1} \frac{B_{N,n-k+1}}{(n-k+1)!} D_x^{n-k},$$

$$(3.14) \quad \hat{P} = D_x.$$

Then (3.13) gives

$$(3.15) \quad \hat{M}B_{N,n}^{(r)}(x) = \left[\left(x - \frac{r}{N+1} \right) - rN \sum_{k=0}^{n-1} \frac{B_{N,n-k+1}}{(n-k+1)!} D_x^{n-k} \right] B_{N,n}^{(r)}(x).$$

(3.3) with $p_n(x) = B_{N,n}^{(r)}(x)$ gives $\hat{M}B_{N,n}^{(r)}(x) = B_{N,n+1}^{(r)}(x)$, thus by Lemma 3.1, (3.15) implies

$$B_{N,n+1}^{(r)}(x) = \left(x - \frac{r}{N+1} \right) B_{N,n}^{(r)}(x) - rN \sum_{k=0}^{n-1} \frac{B_{N,n-k+1}}{(n-k+1)!} \frac{n!}{(n-(n-k))!} B_{N,k}^{(r)}(x).$$

The result now follows from

$$\frac{1}{(n-k+1)!} \frac{n!}{(n-(n-k))!} = \frac{1}{n-k+1} \frac{n!}{k!(n-k)!} = \frac{1}{n-k+1} \binom{n}{k}.$$

□

Proof of the Theorem 1.5. Replacing n by $n+1$ in Theorem 1.8, both sides produces

$$\begin{aligned}
 (3.16) \quad & \frac{1}{rN} B_{N,n}^{(r)}(x) - \left(\frac{x}{rN} - \frac{1}{N(N+1)} \right) B_{N,n-1}^{(r)}(x) \\
 & + \sum_{k=0}^{n-2} \binom{n-1}{k} \frac{B_{N,n-k}}{n-k} B_{N,k}^{(r)}(x) = 0.
 \end{aligned}$$

In Lemma 3.1, denote by $y = B_{N,n}^{(r)}(x)$, and shift indices to produce

$$(3.17) \quad B_{N,k}^{(r)}(x) = \frac{k!}{n!} y^{(n-k)}, \quad k = 0, \dots, n.$$

This gives

$$(3.18) \quad \begin{aligned} & \frac{1}{rN} y - \left(\frac{x}{rN} - \frac{1}{N(N+1)} \right) \frac{1}{n} y' \\ & + \sum_{k=0}^{n-2} \binom{n-1}{k} \frac{B_{N,n-k}}{n-k} \frac{k!}{n!} y^{(n-k)} = 0 \end{aligned}$$

and writing this as

$$(3.19) \quad \frac{1}{rN} y - \frac{1}{n} \left(\frac{x}{rN} - \frac{1}{N(N+1)} \right) y' + \frac{1}{n} \sum_{k=0}^{n-2} B_{N,n-k} \frac{1}{(n-k)!} y^{(n-k)} = 0,$$

produces the result. □

ACKNOWLEDGMENT

This work was supported by the Kyungnam University Foundation Grant, 2015.

REFERENCES

1. P. Appell, *Sur une classe de polynomials*, Ann. Sci. Eco. Norm. Sup. (2) **9** (1880) 119–144 (French).
2. T. Agoh and K. Dilcher, *Convolution identities and lacunary recurrences for Bernoulli numbers*, J. Number Theory **124** (2007), 105–122.
3. A. Byrnes, L. Jiu, V.H. Moll and C. Vignat, *Recursion rules for the hypergeometric zeta function*, arXiv:1305.1892v1 [math.NT] 8 May 2013.
4. K.-W. Chen, *Sums of products of generalized Bernoulli polynomials*, Pac. J. Math. **208** (2003) (1), 39–52.
5. G. Dattoli, H.M. Srivastava and K. Zhukovsky, *Orthogonality properties of the Hermite and related polynomials*, J. Comput. Appl. Math. **182** (2005), no. 1, 165–172.
6. K. Dilcher, *Sums of products of Bernoulli numbers*, J. Number Theory **60** (1996), 23–41.
7. A. Hassen and H.D. Nguyen, *Hypergeometric Bernoulli polynomials and Appell sequences*, Int. J. Number Theory **4** (2008), 767–774.
8. A. Hassen and H.D. Nguyen, *Hypergeometric zeta functions*, Int. J. Number Theory **6** (2010), 99–126.
9. A. Hassen and H.D. Nguyen, *Moments of Hypergeometric Zeta Functions*, J. Algebra, Number Theory: Advances and Applications **7** (2012), No. 2, pp. 109–129.
10. M.X. He and P.E. Ricci, *Differential equation of Appell polynomials via the factorization method*, J. Comput. Appl. Math. **139** (2002) 231–237.
11. F.T. Howard, *A sequence of numbers related to the exponential function*, Duke Math. J. **34** (1967), 599–615.
12. F.T. Howard, *Some sequences of rational numbers related to the exponential function*, Duke Math. J. **34** (1967), 701–716.
13. K. Kamano, *Sums of products of hypergeometric Bernoulli numbers*, J. Number Theory **130** (2010), no. 10, 2259–2271.
14. M.-S. Kim, *A note on sums of products of Bernoulli numbers*, Appl. Math. Lett. **24** (2011) no. 1, 55–61.

15. M.-S. Kim and S. Hu, *Sums of products of Apostol-Bernoulli numbers*, Ramanujan J. **28** (2012), no. 1, 113–123.
16. D.-Q. Lu, *Some properties of Bernoulli polynomials and their generalizations*, Appl. Math. Lett. **24** (2011), no. 5, 746–751.
17. P. Natalini and A. Bernardini, *A generalization of the Bernoulli polynomials*, J. Appl. Math. **3** (2003), 155–163.
18. H.D. Nguyen, *Generalized binomial expansions and Bernoulli polynomials*, Integers **13** (2013) A11. 1–13.
19. H.D. Nguyen and L.G. Cheong, *New convolution identities for hypergeometric Bernoulli polynomials*, J. Number Theory **137** (2014), 201–221.
20. A. Petojević, *A note about the sums of products of Bernoulli numbers*, Novi Sad J. Math. **37** (2007), 123–128.
21. A. Petojević, *New sums of products of Bernoulli numbers*, Integral Transform Spec. Funct. **19** (2008), 105–114.
22. I. M. Sheffer, *A differential equation for Appell polynomials*, Bull. Amer. Math. Soc. **41** (1935), 914–923.

DEPARTMENT OF MATHEMATICS, SOUTH CHINA UNIVERSITY OF TECHNOLOGY,
GUANGZHOU, GUANGDONG 510640, CHINA

E-mail address: hus04@mails.tsinghua.edu.cn

CENTER FOR GENERAL EDUCATION, KYUNGNAM UNIVERSITY, 7(WORYEONG-DONG)
KYUNGNAMDAEHAK-RO, MASANHAPPO-GU, CHANGWON-SI, GYEONGSANGNAM-DO 631-
701, REPUBLIC OF KOREA

E-mail address: mskim@kyungnam.ac.kr